

REPAIR OF WIDE UNILATERAL CLEFT LIP & ALVEOLUS - HOW IS IT DIFFERENT

Prof. Dr. Dr. Srinivas Gosla Reddy

MBBS, MDS, FRCS (Edin.), FDSRCS (Edin), FDSRCS (Eng.), FDSRCPS (Glasg.), PhD

Dr. Rajgopal R. Reddy

MBBS, BDS, FDSRCPS (Glasg.), PhD

Dr. Ashish Fanan M.D.S.

Dr. Avni Pandey M.D.S.

GSR Institute of Craniofacial Surgery,
Hyderabad India



www.craniofacialinstitute.org

GSR Institute of Facial Plastic Surgery



- Non-profit hospital established in 1996
- Dedicated Cleft & Craniofacial Centre of Excellence
- Presently 1,600 cleft and craniofacial surgeries are done every year
- 3 surgeons and 4 fellows with full support team
- More than 30,000 documented cleft & craniofacial surgeries have been performed since 1996
- 600 primary new born cleft children are registered every year



Unilateral Cleft Lip Defect

A 3-Dimensional Problem



Oral

- Discontinuity and mal insertion of Orbicularis oris muscle causing **horizontal and vertical lip length discrepancy**

Nasal

- Deformity of nasal form caused due to mal insertion of Nasalis and other oro-nasal muscles
- **Displacement of septum**

Alveolar

- **Loss of bony support**

Markus, A. F., and Delaire, J. Functional primary closure of cleft lip. Br. J. Oral Maxillofac. Surg. 31: 281, 1993



Unilateral Cleft Lip Defect

Is the morphology of the unilateral cleft lip defect the same in all patients?



Complete Unilateral Cleft Lip



Without Simonart's band (Type I a)

With Simonart's band (Type I b)



Without complete collapse of nasal dome
and ala (Type II a)

With complete collapse of nasal dome and
ala (Type II b)



Complete Unilateral Cleft Lip



Without difference in level of alveolar
ridges (Type III a)

With difference in level of alveolar ridges
(Type III b)



Problems of Wide Clefts

- Differential height of the alveolar segments.
- Variations in the horizontal width of the segments.
- Inward turning of the Cupid's bow towards Columellar base on non cleft side.
- Leading to Severe shortening of skin for Millard rotation.
- Shortening of vertical Height on cleft side and retraction of tissue into the nasal web.
- Collapsed of the nasal dome and severe deviation of nasal septum.





Before primary lip repair (NAM)

Presurgical Nasoalveolar Orthopedic Molding in Primary Correction of the Nose, Lip, and Alveolus of Infants Born With Unilateral and Bilateral Clefts

BARRY H. GRAYSON, DDS
COURT B. CUTTING, M.D.

This addendum to the "State of the Art Dental Treatment of Predental and Infant Patients With Clefts and Craniofacial Anomalies," by Prah-Andersen (*Cleft Palate Craniofac J.* 2000;37:528-532), offers an extended perspective on this controversial subject. This article reviews the role of combined nasal and alveolar (nasopalveolar) molding in the primary correction of the nose, lip, and alveolus of infants born with unilateral and bilateral clefts. The background of presurgical nasoalveolar orthopedic molding, the technique, and the literature are presented. The proposed benefits of treatment from the traditional techniques of presurgical orthopedics have been shown to be unsubstantiated (Kuijpers-Jagtman and Prah, 1996). A close comparison of the proposed benefits of earlier forms of presurgical orthopedics, along with those of the current technique of nasoalveolar molding, is presented.

KEY WORDS: *bilateral unilateral cleft lip and palate, gingivoperiosteoplasty, nasal stent, nasoalveolar molding, nonsurgical columella elongation, presurgical orthopedics*



Presurgical Nasoalveolar Orthopedic Moulding in Primary Correction of the Nose, Lip, and Alveolus of Infants Born with Unilateral and Bilateral Clefts

Dr. Barry H. Grayson, DDS, Dr. Court B. Cutting, M.D. *The Cleft Palate-Craniofacial Journal* Vol 38, Issue 3, pp 193 – 198, May 2001



www.craniofacialinstitute.org

We don't believe in NAM.
Due to **burden of care**.

So

We believe in **Morpho- Functional** Correction.



Goals of Morphofunctional Correction of Unilateral Cleft Lip Defects


A **functional anatomical** repair of the underlying hard and soft tissues is essential.

Goals of primary cleft lip repair

- Harmonious **lip form** in vertical and horizontal dimension
- **Nasal** symmetry
- **Bridging** the alveolar ridge



Millard's Incision for Unilateral Cleft Lip (1996-2000)



Produces better results where

- preoperatively there was a more prominent Cupid's bow and
- where the width of the lip and nostril on the cleft (lateral) side were greater than mean values

Source:


Choice of Incision for Primary Repair of Unilateral Complete Cleft Lip: A Comparative Study of Outcomes in 796 Patients.

Gosla Srinivas Reddy et. al.; Plastic Reconstr. Surg.; 121: 932, 2008



www.craniofacialinstitute.org

Pfeifer's Incision for Unilateral Cleft Lip (2000-2003)



Produces better results

- where the height of the lip on the cleft side was greater and
- where the columella height and width were greater than mean values

Source:

Choice of Incision for Primary Repair of Unilateral Complete Cleft Lip: A Comparative Study of Outcomes in 796 Patients.

Gosla Srinivas Reddy et. al.; Plastic Reconstr. Surg.; 121: 932, 2008



www.craniofacialinstitute.org

Choice of Incision for Primary Repair of Unilateral Complete Cleft Lip: A Comparative Study of Outcomes in 796 Patients

Gosia Srinivas Reddy, B.D.S.,
M.D.S.

Roger M. Webb, F.D.S.,
R.C.S., M.R.C.S.

Ragupathi R. Reddy, B.D.S.

Likith V. Reddy, D.D.S., M.D.

Peter Thomas, B.Sc. (Hons.),

Ph.D.

A. F. Markus, F.D.S.R.C.P.S.,

F.D.S.R.C.P.S.

Hyderabad, India; Plastic
Central Kingston, and Cincinnati, Ohio

Background: No one technique of cleft lip repair consistently produces ideal aesthetic and functional results. This study was carried out in a developing, high-volume center. It compares outcomes attained using two different designs of skin incision used for primary closure of unilateral complete cleft lip and sought to identify the most appropriate technique for clefts of varying morphology.

Methods: Seven hundred ninety-six patients were entered into the study. In each group of slightly less than 400 patients, either a modified Millard or Pfeiffer way line incision was used, both in conjunction with functional repair of the underlying tissues as described by Delaire. Soft-tissue measurements of the lip and nose were recorded preoperatively. Analysis was based on postoperative assessment of the white roll, vermillion border, scar, Cupid's bow, lip length, and nostril symmetry and appearance of the alar dome and base.

Results: Comparison of the two cohorts using Pearson chi-square testing for association and linear trend found a Millard incision gave significantly better results for vermillion match, whereas the Pfeiffer method led to a better postoperative lip length. Preconceptions that one particular technique was better suited to certain preoperative cleft anatomical forms were not proven statistically.

Conclusions: Certain preoperative anatomical features may lead the surgeon to choose one particular incision pattern in preference to another, but in this study, it was found that one technique was essentially as good as the other. This suggests that the technique for closure of the underlying tissues is probably of more importance. (*Plast. Reconstr. Surg.* 121: 932, 2008.)

Surgeons have repaired the deformity of cleft lip for the past 2000 years, since the first attempt performed during the Chin Dynasty in China.¹ Many techniques have been used since that time, and it is clearly apparent that no agreement exists as to which represents the optimum method.

Historically, incisions have been either straight line or broken line, but more recently, in the twentieth century, flap design developed over two distinct periods. In the first, up to 1949, and including Le Mesurier,² lengthening of the lip on the left side was

achieved with some sacrifice of the ipsilateral Cupid's bow. This maneuver, however, tended to produce an aesthetically unfavorable peaking of the lip. In the second half of the century, several attempts were made to counter this shortcoming. Tennison³ utilized a triangular flap on the external surface of the lower margin of the lip, while Pent and Psamme⁴ used a superiorly based flap. Nevertheless, because of scar contracture, this latter approach also produced unacceptable aesthetic outcomes. A combination of superior and inferior flaps was used by Trauner⁵ and Skoog⁶ to counter these problems. A further alternative was described by Malek,⁷ who used a flap based on a precisely measured equilateral triangle to achieve perfect equality in the length of

From the GSR Institute of Craniofacial Surgery, the South Coast Higher Surgical Training Program in Maxillofacial Surgery, the Division of Oral and Maxillofacial Surgery, University of Cincinnati, Dorsal Research and Development Support Unit, Bournemouth University, and Dorsal Cleft Center.

Received for publication March 24, 2006; accepted December 15, 2006.

Copyright ©2008 by the American Society of Plastic Surgeons

DOI: 10.1097/01.prs.0b000200282.05111.4

Disclosure: None of the authors has any financial interest in this work, and no competing interests are declared.

- The Millard flap produced better results when there was a need to rotate the cupid's bow
- Pfeiffer's design produced better results in the vertical elongation of the lip

It was found that one technique was essentially as good as the other.

Choice of Incision for Primary Repair of Unilateral Complete Cleft Lip: A Comparative Study of Outcomes in 796 Patients.
Plastic and Reconstructive Surgery 121: 932, 2008



An incision utilizing the advantages of both Millard and Pfeifer incision

Afroze incision

- Developed to address the problem of **lip length discrepancy and vermillion matching** using only one incision.
- Combined the **Millard incision on the non-cleft side** (medial side) and the **Pfeifer incision on the cleft side** (lateral side).
- **Millard incision** on the non-cleft side **aids rotation** and the **Pfeifer incision** on the cleft side **aids lengthening** trying to address horizontal and vertical discrepancies of the lip.

Source:

Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.





www.craniofacialinstitute.org

Afroze Incision

The Afroze incision **does not cross onto the base of columella.**

Incisions which **cross the** columella **cause scarring** leading to growth retardation and severe downward pull of the columella on affected side



The Afroze incision separates the medial part of ala on cleft side and its associated mal-aligned muscle to further lift the tip of the nose and improve the alar contour and reduce the webbing in the nose

Source:

Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



Afroze Incision for Functional Cheiloseptoplasty

Gosla Srinivas Reddy, DDS, MD,* Rajgopal R. Reddy, BDS, MBBS,* Nilesh Pagaria, BDS, MDS,*
and Stefaan Berge, MD, DD, PhD†

Abstract: Repair of unilateral cleft lip is a fascinating and challenging procedure. Although a great number of operations have been described for the unilateral cleft lip repair, none fulfill all the plastic surgical criteria, and in most cases, cleft lip repairs require secondary operations in an attempt to achieve described goals of primary cheiloplasty. The Afroze incision is a combination 2 incisions, that is, the Millard incision on the noncleft side and Pfeiffer incision on the cleft side. The flap design is the Millard flap on the noncleft side rotated downward, and the peak of the distal curve of the Pfeiffer flap is positioned in the triangular defect formed by the movement of the Millard flap. The proximal curve lengthens downward to receive the Millard's "C" flap. The advantage of this technique is that there is no tension on the postoperative scar because the incision is essentially horizontal in nature, and the contracture of the scar occurs horizontally rather than vertically. Primary septal repositioning is performed, which provides stability and exact positioning of the previously lifted alar crus of the cleft side and nasal tip, and the nose can grow in a balanced way with equal muscular force being exerted on both sides. This incision can be used in all types of complete unilateral cleft lip regardless of the width of the cleft, shortening the cleft lip segment.

Key Words: Complete unilateral cleft lip, Afroze incision, cheiloseptoplasty

J Craniofac Surg 2009;20: 1733-1736

Repair of unilateral cleft lip is a fascinating and challenging procedure. The aims of a unilateral cleft lip repair are to achieve a lip length on the cleft side matching that on the normal side, an inconspicuous residual scar that does not cross anatomic boundaries, an adequate Cupid's bow width, an absence of notching of the vermillion border (whistle-tip deformity), and an absence of peaking of the vermillion at the Cupid's bow on the cleft side. Although a great number of operations have been described for the unilateral

cleft lip repair, none fulfill all the above criteria, and in most cases, cleft lip repairs require secondary operations in an attempt to achieve this described goal.¹

The Millard repair is based on a rotation flap on the noncleft (medial) side coupled with an advancement flap on the cleft (lateral) side. One of its main advantages is that the technique allows adjustment as the operation proceeds, with further rotation and advancement movements tailored to the individual case. It requires the approximation of a pair of convex curves that ultimately may leave a scar crossing the midline at the base of the columella. The Pfeiffer incision is designed using the concept of "morphologic order." Measurements of noncleft side height and length are recorded and translated to the cleft side using a flexible wire, thus determining natural anatomic points. The 2 curves are brought together such that the highest and lowest points of 1 curve are approximated with the corresponding highest and lowest points of the other, thus creating a straight line.²

On comparison of the 2 techniques, each has its own advantages and shortcomings. The Millard flap produced better results when considering vermilion approximation. In this respect, it is rather more flexible than a straight line design, and the operator is able to position the rotation flap on the noncleft side where it is judged likely to produce the best outcome. This technique also has an improved outcome where preoperatively the lip is wider on the noncleft side. This would lead to a reduction in rotational requirement of the flap on the medial side, resulting in less distortion and a Cupid's bow with better form. Repairs using flaps according to Pfeiffer's design resulted in a better length of lip postoperatively. By its nature, the more waves incorporated in the incision, the greater the height of the lip. A prominent wave placed just above the mucocutaneous junction will tend to exaggerate this factor.³

Afroze incision is a combination of 2 incisions. Millard incision on the noncleft side and Pfeiffer incision on the cleft side. The flap design is such that Millard flap on the noncleft side is rotated downward, and the peak of the distal curve of the Pfeiffer flap is positioned in the triangular defect formed by the movement of the Millard flap. The proximal curve lengthens downward to receive the Millard's "C" flap. The advantage of this technique is that there is no tension on the postoperative scar because the incision is essentially horizontal in nature and the contracture of the scar occurs horizontally rather than vertically. There is also no pressure on the Cupid's bow for the same reason.

INCISION MARKING

On the noncleft side, the Cupid's bow is marked by 3 points. Point 1 is the highest point on the contralateral white roll, point 2 is the deepest point on the white roll. Point 3 is marked on the white roll at a distance that is 2 mm more than the distance between points 1 and 2.

On the cleft side, point 4 is marked at a point where the white roll begins to fade (Figs. 1-3).

The Millard incision on the noncleft side is extended from point 3 along the junction of skin and vermillion mucosa and further

From the *DSR Institute of Craniofacial Surgery Hyderabad, Andhra Pradesh, India, and †Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands.
Received January 6, 2009.

Accepted for publication February 28, 2009.

Address correspondence and reprint requests to Gosla Srinivas Reddy, DDS, MD, DSR Institute of Craniofacial Surgery, Postal Address: 17-1-38/3-5, Vinay Nagar Colony, T-5 road, Saidabad, Hyderabad, Andhra Pradesh 500095, India; E-mail: gosla@craniofacialinstitute.org

This article did not require any sources of funding.

The authors declare that they had no financial interests or commercial associations during the course of this study.

Copyright © 2009 by Mutaz B. Habal, MD

ISSN: 1049-2275

DOI: 10.1097/SCS.0b013e3181673a23

The Journal of Craniofacial Surgery • Volume 20, Supplement 2, September 2009

1733

Copyright © 2009 Mutaz B. Habal, MD. Unauthorized reproduction of this article is prohibited.


Afroze Incision for Functional Cheiloplasty,
J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org

Morpho-functional Cleft Lip Repair

Incision design for unilateral cleft lip surgery



Source:

Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org

Morpho-functional Cleft Lip Repair



Minimal muscle dissection on cleft side
ensuring dissection of Orbicularis Oris and **Alar head of
Nasalis muscle**

Source:

Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org

Morpho-functional Cleft Lip Repair



Wide sub-periosteal dissection is done from the vestibule on the cleft side over the piriform rim, nasal bone, infraorbital and malar to lift the **facial mask**

Source:


Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org

Morpho-functional Cleft Lip Repair



Minimal muscle dissection is done on the non-cleft side
relieving all abnormal attachments on anterior nasal spine and
columella

Source:


Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org

Morpho-functional Cleft Lip Repair



SEPTUM IS KEY

The septum is positioned in its rightful anatomical position

Source:


Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org

Morpho-functional Cleft Lip Repair



Perialveoloplasty is done to exert more medial pressure on the palatal shelves

Source:


Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org



At the time of primary lip repair (Morphofunctional Cleft Lip Repair-**Perialveoplasty**)



At the time of lip repair




At the time of palate repair



Morpho-functional repair of complete unilateral cleft lip to achieve aesthetic balance between the lip and nose: an evidence based study Gosla-Reddy, S. et al. International Journal of Oral and Maxillofacial Surgery, Volume 44 , e13 - e14, 2015.



Morpho-functional Cleft Lip Repair



Ala of nose stabilized symmetrically to match that of the normal side by taking a suture through the alar head of the nasalis muscle on the cleft side to the contralateral muscle through the septum

Source:

Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org

Morpho-functional Cleft Lip Repair



OrbicularisOris muscle approximation and closure is done

Source:

Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.



www.craniofacialinstitute.org


Does this incision design protect the
vascularity of the lip?



What we have identified in naso-labial vasculature in cadaver dissection

Morphological and functional variability

- Superior Labial Artery
Caliber asymmetry
- Superior Labial Artery
Anastomosis Inconsistent
- Superior Labial Artery
Duplications
- Philtral Artery
Redundancy Medially
- Philtral Artery
Asymmetry Laterally
- Facialis Artery
Asymmetry




Measurements of S_vO_2 , rHb, flow, (O_2 -metab.) in 2 anatomical planes:

Tissue spectroscopy



0.4 mm → skin

Laser doppler flowmetry



4 mm → muscle




8 surgical landmarks

22 normal




mean age 62m (SD 43)

33 unilat. Cleft
preop



mean age 9m (SD 6)

29 unilat. cleft
Late postop



mean age 23m (SD 48)
time postop 27.5m (SD 33.6m)



Intraoperative Vascular Anatomy, Arterial Blood Flow Velocity, and Microcirculation in Unilateral and Bilateral Cleft Lip Repair

Andreas A. Mueller, M.D.,
D.M.D.

Dietmar Schumann, M.D.,
D.M.D.

Rajagopal R. Reddy,
M.B.B.S., B.D.S.

Katja Schwennner-Zimmerer,
M.D., D.M.D.

Magdalena Müller-Gierth,
M.D.

Hans-Florian Zeitlinger,
M.D., D.M.D.

Hermann F. Söller, M.D.,
D.M.D.

Sevinza Goda Reddy, M.D.,
M.B.B.S., Ph.D.

Read and Smith, Switzerland; and
Hofheim, Germany



Background: Cleft lip repair aims to normalize the disturbed anatomy and function. The authors determined whether normalization of blood circulation is achieved. **Methods:** The authors measured the microcirculatory flow, oxygen saturation, and hemoglobin level in the lip and nose of controls ($n = 22$) and in patients with unilateral and bilateral cleft lip–cleft palate. The authors measured these parameters before lip repair ($n = 29$ and $n = 11$, respectively), at the end of lip repair ($n = 27$ and 10 , respectively), and in the late postoperative period ($n = 33$ and $n = 20$, respectively). The arterial flow velocity was measured in unilateral groups at the same time points ($n = 18$, $n = 11$, and $n = 12$, respectively). Statistical differences were determined using analysis of variance.

Results: Before surgery, the arterial flow velocities and microcirculation values were similar on each side of the face and between groups. The microcirculatory flow was significantly higher in the prelabium of bilateral patients than in the philtrum of controls. All circulation values in unilateral and bilateral patients in the late postoperative period were within the range of controls and of those before surgery. Intraoperatively, the authors consistently found a perfusing artery on the superficial side of the transverse nasalis muscle.

Conclusions: There appears to be no intrinsic circulatory deficit in unilateral and bilateral cleft lip–cleft palate patients. The increased flow in the prelabium indicates a strong hemodynamic need in this territory, compelling to vascular preservation. Whether surgical preservation of the nasalis perforans artery is of long-term benefit should be addressed in future studies. (*Plast. Reconstr. Surg.* 130: 1120, 2012.)

CLINICAL QUESTION/LEVEL OF EVIDENCE: Therapeutic, V.

Cleft lip repair techniques differ mainly in the design of the skin incisions, how the muscle portions are reconstructed, and how the nasal framework is repositioned.¹ The vascular anatomy has remained largely unaddressed in current

surgical techniques, and the reasons for this have yet to be explored.

Normal blood supply is a precondition for development and growth. Thus, it would be of clinical interest to determine whether cleft anatomy leads to a change in the blood supply before or after surgery.

Current techniques for cleft lip repair exclude surgical anastomosis of the lip artery. However, this clinical approach is not based on blood circulation data and so the current standard must be challenged. Vascular damage in cleft surgery interrupts the existent hemodynamics and necessitates further trauma to stop the bleeding, after which the blood circulation may take several months to recover.² Gentle surgical soft-tissue han-

From Craniofacial Surgery, University Hospital Bonn; the HighTech Research Center of Craniofacial Surgery, University of Bonn; the G. R. Institute of Craniofacial and Facial Plastic Surgery; the Anatomical Institute, Neuroscience and Musculoskeletal Anatomy, Laboratory for Functional Morphology; and Cleft-Children International CZL. Received for publication January 17, 2012; accepted May 23, 2012.

Presented in part at the 26th Congress of the European Association for Craniofacial Surgery, in Bruges, Belgium, September 14 through 17, 2010; the 34th International Board-Special-Symposium for Innovative and Vascular Technology in Craniofacial Surgery, in Basel, Switzerland, June 17 through 19, 2010; and the 3th European Craniofacial Congress, in Salzburg, Austria, September 14 through 17, 2011. Copyright © 2012 by the American Society of Plastic Surgeons DOI: 10.1097/PRS.0b013e318267d46b

Disclosure: None of the authors has any conflicts of interest to declare.

Vascular adaption
normal microcirculation late
postoperative in cleft lips.

Columella shows a flow oversupply,
which is maintained late postoperative.

Intraoperative Vascular Anatomy, Arterial Blood Flow Velocity and Microcirculation in Unilateral and Bilateral Cleft Lip Repair
Plastic and Reconstructive Surgery 130 (5): 1120-1129, 2013



Comparison of Three Incisions to Repair Complete Unilateral Cleft Lip

Srinivas Goda Reddy, M.D.S.,
M.B.B.S.

Rajeev R. Reddy, B.D.S.,
M.B.B.S.

Ewald M. Bronckhorst, Ph.D.,
M.D.S.

Anne Marie Kuipers-Jagman,
D.D.S., Ph.D.

Stefaan Berge, M.D., D.D.S.,
Ph.D.

*Hyderabad and Mangalore,
Karnataka, India; and Nijmegen,
The Netherlands*

Background: The incision design for correcting a unilateral cleft lip is important because all subsequent stages of surgery depend on the access and maneuverability of the incision. This prospective cohort study compares the aesthetic and functional outcomes of three different skin incisions for primary unilateral cleft lip repair.

Methods: Patients with complete unilateral cleft lips ($n = 1200$) were enrolled and divided into three groups of 400 patients. Each group of patients was operated on with the Millard incision, Pfeiffer wave line incision, or Afroze incision. Outcome assessments were performed 2 years postoperatively and consisted of assessment of the white roll, vermillion border, scar, Cupid's bow, lip length, nostril symmetry, and appearance of alar dome and base.

Results: With regard to white roll, vermillion border, scar, Cupid's bow, and lip length, the Afroze incision always gave superior results compared with the Millard or Pfeiffer incision. Depending on the cut-off for treatment success, the Afroze incision also showed better results regarding nostril symmetry. With respect to the alar base and alar dome, all three incisions showed comparable outcomes.

Conclusion: The Afroze incision is superior regarding a broad spectrum of outcomes in a heterogeneous population of patients with unilateral cleft lip. (*Plast. Reconstr. Surg.* 125: 1208, 2010.)

The anatomical basis for a cleft lip defect is far removed from the normal orientation. Compared with the noncleft patient, the three groups of superficial facial muscles (i.e., the nasolabial, bilabial, and labiomental) are all displaced inferiorly.¹ The orbicularis oris muscle finds a new and abnormal insertion on the cleft side,² and a partially distorted insertion on the noncleft side.² The Cupid's bow on the cleft side and the white skin roll on both sides are also distorted.⁴ The treatment goals for cleft lip defects are early correction of the cleft, with primary correction to a tension-free, mobile, and balanced lip.¹

The repair of any cleft lip deformity should take not just incision lines into account. A functional anatomical repair of the underlying hard

and soft tissues is essential. Manipulation and repositioning of the microcutaneous tissues must be addressed only once sound foundations have been laid. A primary surgical approach that allows natural facial growth and development, minimizing the need for future secondary procedures, should be every cleft surgeon's goal.²

Many surgical techniques and flap designs have been documented to repair unilateral cleft lips.³⁻¹⁰ Probably the most commonly used is the rotation-advancement technique described by Millard.^{11,12} The Millard incision is based on a rotation flap on the noncleft side coupled with an advancement flap on the cleft side.^{11,12} In our form or another, it is the most widely practiced method today.³

The Pfeiffer incision is designed using the concept of "morphologic order." Measurements of the noncleft side height and length are recorded and translated to the cleft side using a flexible wire, thus determining natural anatomical points.

Disclosure: The authors have no financial interest in this work, and no competing interests are declared.

From the GSK Institute of Craniofacial Surgery, the Department of Preventive and Curative Dentistry, Radboud University Nijmegen Medical Center; A. R. Shetty Memorial Dental College and Hospital, and the Department of Orthodontics and Oral Biology, Cleft Palate Craniofacial Unit, and the Department of Oral and Maxillofacial Surgery, Radboud University Nijmegen Medical Center.

Received for publication June 12, 2009; accepted October 21, 2009.

Copyright ©2010 by the American Society of Plastic Surgeons

DOI: 10.1097/PRS.0b013e3181d15145

1208

www.PRSJournal.com

Copyright © American Society of Plastic Surgeons. Unauthorized reproduction of this article is prohibited.

- Afroze incision performed better
 - Cupids bow position
 - Lip length
 - Lip height
- Millard Incision performed
 - Scar position

What about the nose?

Comparison of Three Incisions to Repair Complete Unilateral Cleft Lip.
Plastic and Reconstructive Surgery, 125 (4): 1208-1216, 2010.



www.craniofacialinstitute.org

Is Primary Septoplasty necessary???

No negative sequelae can be observed after manipulation of the septum in children.


(Smahel, Z. 1999)

Growth of the nose is favorable after primary rhinoplasty.

(McComb, H 1996)



Complete Unilateral Cleft Lip



Without Simonart's band (Type I a)

With Simonart's band (Type I b)



Without complete collapse of nasal dome
and ala (Type II a)

With complete collapse of nasal dome and
ala (Type II b)



Complete Unilateral Cleft Lip



Without difference in level of alveolar
ridges (Type III a)


With difference in level of alveolar ridges
(Type III b)

COMMON FACTOR IN ALL UNILATERAL COMPLETE CLEFT LIPS

DEVIATED NASAL SEPTUM




Is Primary Septoplasty necessary???



A fifteen year old patient with no primary septoplasty



SEPTOCHEILOPLASTY: Unilateral Cleft Lip



- Perichondrium is reflected on both sides of the septum
- The septum is lifted off the nasal spine
- The septum is positioned in its anatomical center
- Perichondrium is closed
- Nasalis muscle from both sides are approximated to form a sling with the septum in the new central position


Source:

Afroze Incision for Functional Cheiloplasty, Technical Note

Gosla Srinivas Reddy et. al.; J. Craniofac. Surg. 20(8):1733-1736, September 2009.





Septocheiloplasty: 1 year post operatively




Septocheiloplasty: 3 years post operatively



Septocheiloplasty: 8 years post operatively



Septocheiloplasty: 15 years post operatively




2 Dimensional Photographic Analysis



Septocheiloplasty: Measuring Outcomes

2 Dimensional Photographic Analysis



Primary Cheiloplasty **without** Septoplasty

Note the septal deviation and alar droop

Source:


Gosla Reddy S, et al. Primary Septoplasty in the Repair of Unilateral Complete Cleft Lip and Palate. Plastic and Reconstructive Surgery, 127 (2): 761-767, 2011



www.craniofacialinstitute.org

Septocheiloplasty: Measuring Outcomes

2 Dimensional Photographic Analysis



Primary Cheiloplasty **with** Septoplasty

Note the absence of septal deviation and reduced alar droop

Source:

Gosla Reddy S, et al. Primary Septoplasty in the Repair of Unilateral Complete Cleft Lip and Palate. Plastic and Reconstructive Surgery, 127 (2): 761-767, 2011



www.craniofacialinstitute.org

Primary Septoplasty in the Repair of Unilateral Complete Cleft Lip and Palate

Srinivas Gosla-Reddy,
M.B.B.S., M.D.S.
Krisztian Nagy, M.D., D.D.S.
Maurice Y. Mommaerts,
M.D., D.M.D., Ph.D.
Rajagopal R. Reddy, M.B.B.S.,
B.D.S.
Ewald M. Bronkhorst, Ph.D.
Rajendra Prasad, B.D.S.,
M.D.S.
Anne Marie Kuipers-Jagman,
D.D.S., Ph.D.
Stefaan J. Berge, M.D., D.D.S.,
Ph.D.

Hyderabad and Mangalore, India;
Bruges-Ostend, Belgium; and
Nijmegen, The Netherlands

Background: The purpose of this study was to assess and compare nasal symmetry in patients who underwent correction of a complete unilateral cleft lip using the Afrozé incision without and with primary septoplasty using a standardized two-dimensional photographic analysis.

Methods: A prospective cohort study of 190 consecutive patients with complete unilateral cleft lip and alveolus with cleft palate treated with or without septoplasty using the Afrozé incision technique was conducted at a high-volume center. Eighty-two patients operated on without primary septoplasty and 76 patients operated on with primary septoplasty were evaluated. Nasal symmetry was compared between patients using two-dimensional photographic analysis. Ratios between the cleft side and the non-cleft side for five parameters were used to assess symmetry: alar base-to-interpupillary line distance, columella-to-Cupid's bow distance, nostril gap area, nostril width, and nostril height. The Mann-Whitney *U*-test was used to calculate differences between the two groups.

Results: Patients operated on with primary septoplasty showed more nasal symmetry compared with patients operated on without septoplasty. This difference was statistically significant for columella-to-Cupid's bow distance, nostril gap area, and nostril height ($p = 0.008$, $p < 0.001$, and $p < 0.001$, respectively) and for the distance between alar base and the alar base-to-interpupillary line distance ($p = 0.145$) the difference was present but not statistically significant. For nostril width, no difference was found ($p = 0.850$).

Conclusion: Patients treated with primary septoplasty showed better results in terms of nasal symmetry when analyzed using two-dimensional photographic analyses. (*Plast. Reconstr. Surg.* 127: 761, 2011.)

Despite a multiplicity of surgical approaches to its correction and as much variation in treatment philosophy, the cleft lip nasal deformity remains a formidable challenge to the reconstructive surgeon treating patients with these congenital deformities. Historically, correction of the cleft nose deformity had been delayed until nasal growth was complete.¹ Early surgical intervention was thought to interfere with normal growth, leading to poor long-term results.¹ Patients with cleft nose deformity had to tolerate the physical nasal deformity and the

psychological trauma well into their adolescence.¹ Randall noted that these patients often were more concerned with their nasal deformity than with their lip deformity.²

Refinement of rhinoplasty techniques has facilitated the ability to address the deformity associated with cleft lip.³ McComb⁴ and Anderl⁵ have published long-term studies that show very little impact on growth with primary correction of the nose deformity along with the correction of the cleft lip. Nevertheless, controversy remains regarding the best time to attempt primary surgical correction of unilateral cleft lip nasal deformity.³⁻⁷ Although a growing number of centers perform the nasal repair in conjunction with cleft lip surgery, some choose a secondary rhinoplasty at a later stage, when the car-

From the GSR Institute of Craniofacial Surgery; Bruges Cleft and Craniofacial Center; Departments of Cariology and Preventive Dentistry, Orthodontics and Oral Biology and Head Cleft Palate Craniofacial Unit, and Oral and Maxillofacial Surgery, Radboud University Nijmegen Medical Center; and A. B. Shetty Memorial Dental College and Hospital. Received for publication July 8, 2010; accepted August 26, 2010.

Copyright ©2011 by the American Society of Plastic Surgeons
DOI: 10.1097/PRS.0b013e318200a97a

Disclosure: None of the authors has any financial interest in this work, and they have no competing interests to declare.

Primary septoplasty showed better results in terms of nasal symmetry when analyzed using two-dimensional photographic analyses.

Primary Septoplasty in the Repair of Unilateral Complete Cleft Lip and Palate.
Plastic and Reconstructive Surgery, 127 (2): 761-767, 2011



3 Dimensional Photographic Analysis





3 Dimensional Photographic
Equipment




3 Dimensional LASER
Equipment



Measurement: Right Nostril (Transversal)



Right Nostril Transversal: 12.1 mm



Right Nostril Transversal: 12.9 mm

3D Stereophotogrammetric analysis supported by **Radboud University, Nijmegen (Prof. Stefaan Berge)** and **University Medical Center, Basel (Prof. Hans Florian Zeilhofer)**



www.craniofacialinstitute.org

Landmarks & Measurements

3 D Photographs and LASER Images



Results

3 Dimensional Nasal Analysis of Patients with Complete Unilateral Cleft Lip corrected with Septocheiloplasty

Volumetric analysis of the nose



Source:

Gosla Reddy et.al. 3D Stereo photo grammetric analysis of lip and nasal symmetry after primary cheiloseptoplasty in primary cleft lip repair.

Rhinology, 49: 546-553, 2011





www.craniofacialinstitute.org

Results



3 Dimensional Nasal Analysis of Patients with Complete Unilateral Cleft Lip corrected with Septocheiloplasty

Transverse/Horizontal Nostril Length



Mean Symmetry ratio of 1.25

Vertical Nostril Length



Mean Symmetry ratio of 0.97

Source:

3 Dimensional Analysis of Patients with Complete Unilateral Cleft Lip corrected with Septocheiloplasty.


Gosla Reddy S, Mommaerts MY, Reddy R, Chaitidis D, Mueller A, Schwenzer K, Berge S: Ongoing Study, Radboud University, Netherlands and University of Basel, Switzerland



Results

3 Dimensional Nasal Analysis of Patients with Complete Unilateral Cleft Lip corrected with Septocheiloplasty

Volumetric analysis of the nose



Ratio Left Volume vs. Right Volume = 1.09

Source:

Gosla Reddy et.al. 3D Stereophotogrammetric analysis of lip and nasal symmetry after primary cheiloseptoplasty in primary cleft lip repair.

Rhinology, 49: 546-553, 2011



www.craniofacialinstitute.org

3D stereophotogrammetric analysis of lip and nasal symmetry after primary cheiloseptoplasty in complete unilateral cleft lip repair*

Bram van Loon^{1,3,4}, Srinivas G. Reddy^{2,3}, Niels van Heerbeek^{3,4}, Koen J.A.O. Ingels^{1,4}, Thomas J.J. Maal^{1,4}, Wilfred A. Borstlap^{1,4}, Rajgopal R. Reddy², Anne-Marie Kuijpers-Jagtman³, Stefaan J. Bergé^{1,4}

¹ Department of Oral and Maxillofacial Surgery, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

² GSR Institute of Craniofacial Surgery, Hyderabad, India

³ Department of Otorhinolaryngology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

⁴ Facial Imaging Research Group, Nijmegen, Bruges

⁵ Department of Orthodontics and Oral Biology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

All authors are united in the Centre for Facial Plastic Surgery, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

SUMMARY

Background: The aim of this study was to evaluate symmetry of the lip and nose in patients with CUCLP after primary cheiloseptoplasty (Afroze technique), in comparison to non-cleft controls.

Methodology: In this prospective study, forty-four patients with operated non-syndromic CUCLP were included. The control group consisted of 44 volunteers without cleft defects of approximately the same age and sex. Primary septoplasty was performed in conjunction with the cleft lip (CL) repair using the Afroze incision. 3D facial images were acquired using 3D stereophotogrammetry. After a 3D cephalometric analysis of the lip and nose was performed in both groups, linear and volumetric data were acquired. Lip and nose symmetry were calculated and compared using Student's t-tests as well as the Chi square test.

Results: For all measurements, the control group was up to 36% closer to perfect symmetry compared to the CUCLP group after primary surgery. This difference was statistically significant.

Conclusions: After primary cheiloseptoplasty according to the Afroze technique in patients with CUCLP, asymmetry in the nose and lip area still exists as compared to non-cleft controls. Although non-cleft individuals also show some degree of asymmetry, the results of this study stress the difficulty in obtaining near normal symmetrical relations.

Key words: cleft palate, three-dimensional imaging, maxillofacial surgery, nose, rhinoplasty, 3D stereophotogrammetry, volume.

INTRODUCTION

The ultimate goal for repair of the complete unilateral cleft lip, alveolus and palate (CUCLP) deformity is to create normal oronasal form and function. This aim has resulted in a plethora of techniques and innovations to optimize the esthetic and functional results. However, the management of CUCLP deformities, especially that of the nose, remains a challenge.

Various studies¹⁻⁶ have been undertaken to evaluate the results of different operative procedures to correct the CUCLP nose deformity. However, quantification of rhinoplastic procedures remains difficult. Besides direct anthropometric measurements^{1,2}, studies comparing pre- and postoperative nose and lip changes in patients with clefts are limited to two dimension-

Footnote: #Both authors contributed equally to the study

*Received for publication: May 2, 2011; accepted: August 21, 2011

DOI:10.4193/Rhino.11.092


Primary septoplasty showed better results in terms of nasal symmetry when analyzed using three-dimensional photographic analyses.

3D Stereophotogrammetric analysis of lip and nasal symmetry after primary cheiloseptoplasty in primary cleft lip repair.
Rhinology, 49: 546-553, 2011



My Opinion

The cleft lip defect is a 3 dimensional problem



Only a MorphoFunctional approach that addresses all three dimensions will positively effect the repair of the Unilateral Lip.

My solution

CHEILOPLASTY, SEPTOPLASTY and PERIOPLASTY



Bring the Smile Back

